

Citation for published version:

Malli, M & Forrester-Jones, R 2021, *The experience of stigma among adults with Tourette's syndrome: Executive Summary*.

Publication date:
2021

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



The experience of stigma among adults with Tourette's syndrome.

Executive Summary

Melina Malli and Rachel Forrester-Jones

Research commissioned and funded by Tourettes Action

University of
Kent



Funding

This publication arises from research commissioned and funded by Tourettes Action under Grant 2955. The analyses and report were conducted by the authors. The facts presented and views expressed in this report are those of the authors and not necessarily those of the funders.

Contact information:

Melina Malli, Qualitative Research Associate, Nuffield Department of Primary Care Health Sciences, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG melina.mali@phc.ox.ac.uk

Rachel Forrester-Jones, Professor of Social Policy, Director of the Centre for Analysis of Social Policy, Social Policy and Sciences, University of Bath, Claverton Down, Bath BA2 7AY rfj23@bath.ac.uk

[Cover image by Luis Quintero on Pexels](#)

© Melina Malli and Rachel Forrester-Jones March 2021

All rights reserved. Reproduction of this report by photocopying or electronic means for non-commercial purposes is permitted. Otherwise, no part of this report may be reproduced, adapted, stored in a retrieval system or transmitted by any means, electronic, mechanical, photocopying, or otherwise without the prior written permission of the author.

Acknowledgements

We would like to thank Tourettes Action for advertising this study and their support in the recruitment of participants. Most of all we thank all those with Tourette's syndrome, for their kindness and willingness to talk to us about their life experiences -it has been an honour to work with you all.

“Never laugh at other people's disabilities. Unless they have Tourette's, because how can you not?”

Executive Summary

Stigma devalues individuals depriving them of full social acceptance (Goffman, 1963). Yet despite growing awareness of the negative consequences of stigma for people with disabilities (see Corrigan, Patrick & Bink, 2005; Liamputtong, 2013; Tang & Bie, 2016) our study is one of the first to systematically examine the experience of stigma from the perspective of adults with Tourette's syndrome.

A neurological condition characterised by tics (involuntary unwanted movements and/or vocalisations that can vary in frequency, complexity and intensity (American Psychiatric Association, 2013), TS is often accompanied by obsessive-compulsive disorder (OCD), attention deficit hyperactivity disorder (ADHD), anxiety disorders, and depression. Despite popular stereotyping of the condition as the 'swearing disease' (see Calder-Sprackman, Sutherland, & Doja, 2014) by virtue of coprolalia (involuntary utterances of obscene words), this symptom of TS is fairly uncommon (Freeman et al., 2009). Whilst some progress has been made to de-emphasise this aspect of the condition (see documentaries such as 'Tourette De France' and 'Twitch') TS is still generally portrayed in the media as synonymous with uncontrollable cursing and unpredictable behaviour (Malli, Forrester-Jones, & Murphy, 2016).

The aims of our UK study were:

1. to assess the extent and nature of stigma as experienced by people with TS;
2. to understand how stigma impacted the quality of life of adults with TS;
3. to gain coping strategies adopted by individuals with TS to cope with stigma.

Our overall objective was to inform current policy and practice around TS.

We used a mixed-methods study design. Tourettes Action support groups and an advisory group including individuals with TS and their family carers helped design the study. Participants were recruited via a video <https://www.youtube.com/watch?v=XVQTO0wew8A> posted on relevant platforms (e.g. Tourettes Action, Facebook, Twitter) and through the Research Participation Registry of Tourettes Action database as well as snowball sampling. The study took place between from 2019 to 2020 and was given a favourable opinion by a University of Kent Ethics Committee (REF: 11/2018).

In addition to digitally recorded face-to-face on-line interviews, an on-line survey included the following measures:

- *Discrimination and Stigma Scale (DISC-12)* - assessing the degree to which people with TS experienced stigma and discrimination in their everyday lives, and
- *Perceived Quality of Life* (Patrick et al., 1998; 2000) - to assess individuals' subjective evaluation of their quality of life.

Quantitative data was subjected to statistical analysis and qualitative data was analysed using thematic analysis.

Sample characteristics

199 adults with TS completed the survey. The mean age of participants was 31 (age range 18-73). 48% were female, 46% were male, and 6% identified as "other gender". Over half the sample identified as White British (67%) and the majority (n=125) were single. The majority (56.2%) reported a tic onset between the ages of four to eight, whilst a small number reported an adult onset. Additional (one or more) mental health conditions or developmental disorders were reported by 76.6% of the participants, including Anxiety Disorder, OCD, ADHD and Autism.

20 adults (14 men and 6 women with a mean age of 33 (range: 20-71) self-identifying as having Tourette's were interviewed. 15 reported comorbid diagnoses, including OCD, anxiety, autism, ADHD, dependency, eating disorders.

Quantitative Findings

- Participants reported experiencing discrimination in numerous life domains and most prominently in education (75.7%), interpersonal relationships (71.4%), and employment (54.3%).
- Stigma also appeared to play an important role in social isolation.
- Most participants tried to conceal their condition (75%) and over half had not applied for jobs (54%) or higher education courses (50.3%) for fear of unjust treatment.
- Individuals with comorbidities reported significantly more stigmatisation than those without any co-occurring conditions.
- Experiences of stigma impacted their quality of life significantly and poorer quality of life was also significantly associated with anticipated discrimination. Finally, experienced and anticipated stigma were positively correlated.

Qualitative Findings

- humour that trivialises and reduces TS to simplistic notions mediates the construction and perpetuation of stigma; trivialisation in particular impedes the general public from understanding the complexity of TS;
- this leads to both subtle and overt discriminatory practice in micro-interactions and structural and institutional levels;
- people with TS attempt to manage their 'stigmatised identity' through concealment and self-stigma.

Conclusion

Stigma is an issue of substantial concern for people adults with TS. The experience of TS stigma dramatically impairs quality of life, manifests in different contexts and effects self-worth.

Recommendations

- There is a need for public initiatives that counteract stigma on different levels;
- Practitioners need to pay greater attention to how people with TS try to manage stigma;
- Future anti-stigma interventions need to consider incorporating humour as an educational strategy that will simultaneously entertain but also allow the audience to reflect on their prejudice and stigmatising behaviours;
- More studies that assess the effectiveness of TS anti-stigma interventions on specific community groups (i.e. healthcare professionals, secondary school children, college students, employers) are needed.

Society's collective response to TS need to change as barriers and restrictions still stem from prejudicial attitudes.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (DSM-5®)* American Psychiatric Pub.
- Calder-Sprackman, S., Sutherland, S., & Doja, A. (2014). The portrayal of tourette syndrome in film and television. *Canadian Journal of Neurological Sciences*, 41(2), 226-232.
- Freeman RD, Zinner SH, MÜLLER-VAHL KR, et al. Coprophenomena in tourette syndrome. *Developmental Medicine & Child Neurology*. 2009;51(3):218-227.
- Goffman E. Stigma: Notes on the. *Management of Spoiled Identity*. London: Penguin. 1963.
- Liamputtong P. *Stigma, discrimination and living with HIV/AIDS: A cross-cultural perspective*. Springer Science & Business Media; 2013.
- Malli, M. A., Forrester-Jones, R., & Murphy, G. (2016). Stigma in youth with tourette's syndrome: A systematic review and synthesis. *European Child & Adolescent Psychiatry*, 25(2), 127-139.
- Patrick DL, Kinne S, Engelberg RA, Pearlman RA. Functional status and perceived quality of life in adults with and without chronic conditions. *J Clin Epidemiol*. 2000; 53(8):779-785.
- Patrick DL, Danis M, Southerland LI, Hong G. Quality of life following intensive care. *Journal of general internal medicine*. 1988; 3(3):218-223.
- Tang, L., & Bie, B. (2016). The stigma of autism in china: An analysis of newspaper portrayals of autism between 2003 and 2012. *Health Communication*, 31(4), 445-452.